

Predatory Arthropods For Brown Planthopper Species (*Nilaparvata lugens* Stal): A Comparisional Study Based On Agroforestric

by Dadi Unigal

Submission date: 29-Nov-2021 08:01AM (UTC+0700)

Submission ID: 1714562352

File name: Final_Paper_Scopus_Q3_NVEO.pdf (296.77K)

Word count: 6730

Character count: 36721

Predatory Arthropods For Brown Planthopper Species (*Nilaparvata lugens* Stal): A Comparisonal Study Based On Agroforestric

Dadi Dadi

Universitas Galuh Ciamis-Jawa Barat-Indonesia

*Corresponding author. Email: dadi@unigal.ac.id

Submitted: 27.01.2021; Accepted: 24.10.2021

Abstract

This study describes the potential of agroforestry ecosystems in supporting the diversity of predatory arthropods in rice fields during the planting interval, problems are often found in the planting interval process, so the cause of the error must be found. Methodology: Two locations of rice fields around agroforestry with different types of agroforestry were selected, namely rice fields adjacent to simple agroforestry and complex agroforestry in Gunungcupu Village, Sindangkasih District, Ciamis Regency, West Java. Arthropod species data were obtained using a mask and suctioning was carried out with an aspirator. There are 40 sample plots at each location. Plots were placed systematically in rice fields adjacent to agroforestry with a distance between plots of 10 m and a distance between plots of 10 m. Observations were made since the beginning of the first planting period, September 20, 2018, continued during the planting interval and the end of the planting interval in the last week of the second planting period on June 14, 2019. Results: Based on the results of the analysis, it was found that the even distribution of predatory arthropod species in the rice fields was not affected by the planting period. The planting interval also did not affect the diversity and even distribution of insect predatory arthropod species. This condition indicates that agroforestry areas have the potential to support the diversity of arthropod species in paddy fields.

Keywords: Ecosystem, agroforestry, Planting Break

Introduction

Agriculture has the potential to cause ecological problems. The problems found are mainly related to the imbalance of the system chain in the land which is indicated by the emergence of pests in abundance at certain times. These findings are strengthened after intensive farming with the use of synthetic fertilizers and poisons. Planthoppers in rice fields, for example, have long been an important pest in Japan and other tropical Asian countries (Hattori, 2001). The rice planthopper is endemic to the mainland tropics, but its habitat has temporarily spread to Japan and Korea in a massive long-distance migration (Li et al., 2011; Liu & Lou, 2018; Supriyadi & Wijayanti, 2018). The increasing influence of the brown planthopper (*Nilaparvata lugens* Stal), for example, has occurred since 1970 as a pest that destroys rice harvests in Indonesia, especially in Java, North Sumatra and Bali (Nuryanti et al., 2018; Deosi et al., 2020). In 1976-1977 the area of rice fields attacked by brown planthoppers reached 347,000 ha, thus disrupting the national supply of planting breaks. The brown planthopper attack in waves from time to time is influenced by its physiological properties that are able to create new biotypes to break rice resistance in varieties grown in the field (Wang et al., 2018; Rizkiani & Herlinda, 2018; Dadi, 2019). Green leafhoppers (*Niphotettix virescens*) are also widely reported to be the cause of crop failure (Desiska et al., 2019) mainly because of their ability to

transmit tungro virus which is very damaging to rice crops. This problem is of course closely related to the surrounding environmental conditions. thus disrupting the supply of planting breaks nationally. The brown planthopper attack in waves from time to time is influenced by its physiological properties that are able to create new biotypes to break rice resistance in varieties grown in the field (Isrin & Fauzan, 2019; Srinivasa et al., 2020; Roy et al., 2020). Green leafhoppers (*Niphotettix virescens*) are also widely reported to be the cause of crop failure (Silva et al., 2021) mainly because of their ability to transmit tungro virus which is very damaging to rice crops. This problem is of course closely related to the surrounding environmental conditions. thus disrupting the supply of planting breaks nationally. The brown planthopper attack in waves from time to time is influenced by its physiological properties that are able to create new biotypes to break rice resistance in varieties grown in the field (Nasral et al., 2020). Green leafhoppers (*Niphotettix virescens*) are also widely reported to be the cause of crop failure (hanif et al., 2019; Dadi, 2020) mainly because of their ability to transmit tungro virus which is very damaging to rice crops. This problem is of course closely related to the surrounding environmental conditions. The brown planthopper attack in waves from time to time is influenced by its physiological properties that are able to create new biotypes to break rice resistance in varieties grown in the field (Kardinan et al., 2019; Muazam & Nugroho, 2020). Green leafhoppers (*Niphotettix virescens*) are also widely reported to be the cause of crop failure (Ngatimin et al., 2020) mainly because of their ability to transmit tungro virus which is very damaging to rice crops. This problem is of course closely related to the surrounding environmental conditions. The brown planthopper attack in waves from time to time is influenced by its physiological properties that are able to create new biotypes to break rice resistance in varieties grown in the field (Aliffah et al., 2020). Green leafhoppers (*Niphotettix virescens*) are also widely reported to be the cause of crop failure (Dharshini & Gowda, 2014; Tanaka et al., 2015) mainly because of their ability to transmit tungro virus which is very damaging to rice crops. This problem is of course closely related to the surrounding environmental conditions.

An ecological approach can overcome pest attacks on agricultural land. The approach taken is to present natural enemies of pests on the land (Budiarti et al., 2021). The ecological principle emphasizes that pests and natural enemies must stop planting in a balanced state. Pests do not have a detrimental effect on crops because ecosystems provide natural mechanisms through numerical responses and functional responses. The mechanism is formed through the diet eaten between predators/parasitoids and prey/hosts.

Arthropods have an important role in the ecosystem. In relation to the agricultural ecosystem, there are arthropods that play a neutral role, act as pests and some act as natural enemies of the pests concerned (Sun et al., 2018; Adnan, 2019). Mahrub (1998) conducted a study on the community structure of arthropods in lowland rice in the rainy and dry seasons without the application of insecticides. An important finding from this research is that in the lowland rice ecosystem the diversity and abundance of arthropods is known to be quite high. Pest population is lower than predators. This condition has encouraged negative feedback interactions and can increase the potential for natural enemies so that the natural pest control process runs well without the use of insecticides. Increasing the diversity of habitats in agricultural landscapes can increase the diversity of insect pests and beneficial insects (natural enemies) and often reduce crop damage by pests (Aldini et al., 2019). The creation of such a balance condition raises the next challenge, namely the application of the most suitable strategy in the cultivation area. Consideration to create the right habitat for natural enemies is now important.

Agroforestry can be an alternative habitat for natural enemies. Agroforestry has a main function that emphasizes meeting the needs of life in the form of agricultural production (improvement of welfare) and ecosystem sustainability, from a practical point of view. The ability of agroforestry to maintain and maintain natural resources has an impact on various things, namely: 1) maintaining physical properties and soil sustainability, 2) maintaining the hydrological function of the area, 3) maintaining carbon stocks, 4) reducing greenhouse gas emissions and 5) maintaining biodiversity (Meilin, 2017; Dhaliwal et al., 2019). Agroforestry makes beneficial contributions both economically and ecologically (Widianto et al., 2003). Agroforestry ecosystems can also be a habitat for useful animals such as pollinators and predators of agricultural pests. This is an important value in agroforestry areas related to the condition of the rice field ecosystem. Agroforestry that functions as a forest area is a supporting force for ecosystems that must be maintained for its sustainability. Of conceptual relevance between agroforestry and forests, Zeitoun et al (2010) view forests as important controllers of natural insect diversity. Consideration of forest as a balancer for lowland rice farming ecosystems because rice fields have a fragile ecosystem. The chain system in the rice field ecosystem has been modified to be simpler and more lame (Zeitoun et al., 2016). The agroforestry system approach to provide carrying capacity for lowland rice cultivation is very important.

This picture is interesting to be studied further, especially with regard to the functioning of the agroforestry ecosystem in providing resources for the predatory arthropods of leafhoppers in the rice field ecosystem. The findings of this study are useful for various aspects, both scientific and applied. Practitioners in the field also received new information about the potential for the application of agriculture through insect management systems on agricultural land.

Research location and time

This research is in the form of field exploration studies in farmers' fields. The rice fields chosen as the research location are those in the vicinity bordering the agroforestry area. The study location is in Gunungcupu Village, Sindangkasih District, Ciamis Regency, West Java Province. Ciamis District.

The research location was determined in two rice fields that have different types of agroforestry areas, namely simple agroforestry and complex agroforestry. The rice fields were selected by cooperative farmers who grow rice conventionally, namely farming using inorganic fertilizers and using pesticides for pest control. Data collection in the field was carried out for 2 uninterrupted planting seasons starting on September 20, 2018 until June 14, 2019.

Research tools and materials

The tools needed for transect activities are a plastic mine measuring 200 m long, a lid with a diameter of 55 cm² (Mahrub, 1998; Dadi, 2020), a tape meter, and red bamboo poles with a length of 1 m which are used as markers. permanent sample plot.

The main tools for observing arthropods are the mask and aspirator. The lid is made of a frame made of iron plate covered with mica plastic in the form of a tube with a bottom surface diameter of 55 cm and a height of 80 cm. The aspirator is the result of a modification of a mini vacuum cleaner with energy supply from a 1000 watt mini generator. Another tool needed is a cloth bag to hold the sample in the aspirator. Species identification is carried out in the laboratory, the equipment needed in the field is a drum, sasag, newsprint, soft-bristled brush, plastic bag clip, small styrofoam cubes, label

paper, species recapitulation form, and stationery.

Research methods

This research is a quantitative exploratory study. The research variables included the diversity of plant species in the agroforestry area, the diversity and abundance of predatory arthropod species in the agroforestry and paddy fields. The diversity of plant species in the agroforestry area is used as the basis for comparing the types of agroforestry in the research location. Abiotic environmental conditions were also taken as complementary data including temperature, humidity and daily rainfall.

Data collection in the agroforestry area includes vegetation and arthropods. Vegetation data is about the structure and diversity of plants, while arthropod data is the diversity and evenness of arthropods which are natural enemies of rice planthoppers. The direction of the transect was determined across the agroforestry area bordering the rice field area along 200 m. The diversity and abundance of natural enemy arthropods was carried out using a systematic sampling system.

Observations of predatory arthropods were carried out in rice fields. The number of observation plots is 60 points which are divided into two separate blocks perpendicular to the boundary line of rice fields-agroforestry.

The unit of sample in the fields is the square. The area of the square in the form of a circle is 0.24 m² with a diameter of 0.55 m. Since the object of observation is a moving animal, a square shaped hood is used (Rasul et al., 2005;) taking into account the spacing of rice in the field. The sample plot becomes a permanent observation area regarding the diversity and even distribution of predatory arthropod species. The length of the observation path per block is 30 m towards the rice fields. The decision on line length refers to a study on the effect of weed rows on the agroecology of farmland in Switzerland which found that from an ecological point of view the best distance between weed rows was 24 m (Grimaud-Hervé, et al., 2016).

The distance between the axis points of the sample plots is 3 m. The number of observations per lane is 10 plots. The number of observation lines per area is 3 and the distance between the observation lines is 4 m. Sampling was carried out at intervals of once a week. Observation time starts at 06.00 until 10.30 WIB.

The presence of arthropods was analyzed according to the planting period to provide an overview of the relationship between the growing season and population dynamics. The analysis consisted of three observation phases, namely the analysis of the first planting period, the planting interval and the second planting period. The first planting period was observed on Monday, November 24, 2008 (first week of observation or 8th day after planting) and the last planting interval was on Monday, March 2, 2009 (15th week of observation or the first Monday after harvest). The second phase of observation was carried out during the planting interval, when the rice fields were left by the farmers, empty without cultivated plants. Observations of the planting interval were carried out starting from the 16th week to the 23rd week of observation.

The third phase of observation is during the second planting period. Observations of the second planting period were carried out for 15 weeks starting Monday March 14 (24th week of observation) and ending on Monday 17 June 2019, which is the 38th week or the first Monday after harvest. Chart 1 depicts the observation phase during the research.

Chart 1. Observation phase at the research site

24 Sept – 31 Dec 2018	7 Jan – 25 Feb 2019	March 4 – June 17, 2019
First planting time	Planting break	Second planting time
(15 weeks of observation)	(8 weeks of observation)	(15 weeks of observation)

Samples of arthropod species from each observation plot were identified using identification clues from Sugiyama et al (2006), Rightmire et al (2019) and matched the object of observation with pictures from Sheppard (1995) Samples were identified by species and number of individuals.

Data analysis technique

Data analysis was carried out in two stages, namely index calculation and descriptive analysis. The first step is to calculate the species diversity index and evenness index of arthropod predatory planthopper species in each observation plot. The species diversity index and evenness index of predatory arthropods were calculated using Microsoft Office Excel 2010 software. The species diversity index and evenness index were calculated using the species diversity formula from Shannon-Wiener and the evenness index formula derived from Shannon-Wiener as follows (Heong et al. , 1991; Cheng, 1995):

$$H' = -\sum \left[\left(\frac{n_i}{N} \right) \ln \left(\frac{n_i}{N} \right) \right]$$

$$E = \frac{H'}{\ln S}$$

Information:

- H' = Shannon-Wiener Indeks Index
- n_i = Number of individual species -i
- N = Total number of individuals from the sample plot
- E = Evenness index
- S = Number of all species on the plot

The second stage is the descriptive analysis process. The analysis aims to test the hypothesis proposed in this study. The calculation process is carried out with the help of Microsoft Office Excel 2010 software. At this stage the results of the analysis of species diversity and evenness are interpreted and described. The results of each interpretation are reinforced with graphs to visually illustrate the results of the analysis.

RESULTS AND DISCUSSION

Plant samples were taken using stratified plots. Observations of plant diversity were carried out in two different locations, namely Location A and Location B. The number of samples taken at each location took into account the broad representation of each agroforestry. Based on the results of field measurements, it is known that the area of Location A is 31,095.21 m² and the area of Location B is 47,684.34 m².

The characteristics of Location A are rice fields side by side with simple agroforestry. This agroforestry is dominated by mahogany (*Swietenia mahagoni*). Other plants found with rare frequency were coconut (*Cocos nucifera*), albasiah (*Albizia falcataria*) and banana (*Musa sp.*) while those found very rarely were teak (*Tectona grandis*). The grass plants in the middle of the forest are not too dense because mahagoni as the main plant making up the forest vegetation has a fairly high level of area cover so that the grass plants do not have the opportunity to grow freely. Grass plants that are denser are found on the outskirts of agroforestry bordering the rice field ecosystem. The grasses that grow there are *Laersia sp.*, *Paspalum conjugatum*, *Imperata cylindrica*, *Ageratum conyzoides*, *Phyllanthus niruri*. This grass looks very tight, especially during the rainy season. The cropping pattern on this land does not follow certain rules. *S. mahagoni* as the main constituent of vegetation grows without certain rules.

Ripe seeds that fall around the forest floor then grow naturally so that this plant has a complete strata starting from the new seedling level, saplings, poles to the shape of an adult tree with a trunk diameter varying between 1-27 cm. These forest characteristics are categorized as simple agroforestry. Ripe seeds that fall around the forest floor then grow naturally so that this plant has a complete strata starting from the new seedling level, saplings, poles to the shape of an adult tree with a trunk diameter varying between 1-27 cm. These forest characteristics are categorized as simple agroforestry. Ripe seeds that fall around the forest floor then grow naturally so that this plant has a complete strata starting from the new seedling level, saplings, poles to the shape of an adult tree with a trunk diameter varying between 1-27 cm. These forest characteristics are categorized as simple agroforestry.

The characteristics of Location B are rice fields adjacent to complex agroforestry. The agroforestry at Location B has much more intermittent vegetation. The agroforestry land in this location is managed by the owner by deliberately planting various types of plants without setting certain rows or rows. Similar plants are planted anywhere on land that allows them to be planted. The distances and rows/rows are not well defined so that a regular crop pattern does not appear. Plants consist of seasonal plants, namely kencur (*Kaempferia galangal*), taro (*Colocasia esculenta*), cayenne pepper (*Capsicum frutescens*), turmeric (*Curcuma longa*) and cassava (*Manihot utilissima*). The fruit plants found consisted of banana (*Musa sp.*), jackfruit (*Artocarpus integra*), rambutan (*Nephelium lappaceum*), salak (*Zalacca edule*), randu (*Bombax malabaricum*), cloves (*Eugenia aromatic*), coconut (*C. nucifera*), mango (*Mangifera indica*) and jatropha (*Jatropha curcas*). Timber plants consist of teak (*T. grandis*), albasia (*A. falcataria*), tissuek / mountain waru (*Hibiscus macrophyllus*) and African wood (*Maesopsis eminii*). Land owners use land with any useful plants to meet short, medium and long term needs. Grass plants are allowed to grow to compete with seasonal plants that are deliberately planted. These forest characteristics are categorized into complex agroforestry groups. Land owners use land with any useful plants to meet short, medium and long term needs. Grass plants are allowed to grow to compete with seasonal plants that are deliberately planted. These forest characteristics are categorized into complex agroforestry groups. Land owners use land with any useful plants to meet short, medium and long term needs. Grass plants are allowed to grow to compete with seasonal plants that are deliberately planted. These forest characteristics are categorized into complex agroforestry groups.

Diversity and evenness of arthropod species by location and season

³ Based on the results of the analysis, it is known that the rice fields around which there is a simple type

of agroforestry (Location A) have a lower total number of individual arthropod species predators of rice planthoppers than the rice field where there is a complex type of agroforestry around them (Location B). However, in terms of species diversity, research sites with simple type agroforestry have a higher species diversity index than those in complex type agroforestry research sites.

The type of agroforestry is related to the ¹number of predatory ¹species in the rice fields. Based on the observations, rice fields surrounding complex type agroforestry have more arthropod species than rice fields surrounding simple type agroforestry. ¹Table 1 presents the number of arthropod species predators ¹of the rice planthopper by location and observation phase. ¹The number of predatory arthropod species in rice fields where there is agroforestry is 26 while the number of species in rice fields where there is complex agroforestry is 29. The planting period, on the other hand, does not appear to be related to the interplay of species in the field because the data obtained inconsistent information. from one observation location to another. For example, in Location A, 25 predatory arthropod species were found in the first planting period and at the second planting period, more species were found (26). The findings at Location B were the opposite, because the number of species found in the first planting period was actually more (29 species) than the second planting period (27 species).

The number of predatory arthropod species during the planting interval is less. The data at Location A and Location B showed that during the planting interval the number of predatory arthropod species was much less than the number of species at the planting period, but during the planting interval the number of species at Location A became more (22 species) than Location B (21 species).

The total number of predatory arthropod individuals during the observation was 5,178. The number by location during the observation, individuals at Location B (2,966) was more than Location A (2,122). Thus, the predatory arthropods of rice planthoppers in rice fields where there are complex types of agroforestry ecosystems have 34% more than other locations in this study. The number of individual predatory arthropods at the planting interval at Location B was also much higher (632) than the number at Location A (440), although the number of species showed the opposite (21 and 22 species, respectively). As the description of the number of species, it turns out that the number of individuals found in each research location also does not show the same conditions in the two growing seasons. The highest number of individuals in Location A (1. 027) was found during the second planting season, while in Location B the highest number of individuals (1,586) was found during the first planting season. The determining factor for this condition is probably related to the condition of the paddy fields. It is known, in the second planting period where rainfall has decreased quite sharply causing the condition of the paddy fields to become drier, but specifically at Location A the condition of the paddy fields during the second planting period remains at a planting interval so that it has higher humidity around the planting. This condition probably caused the predatory arthropod community to be more abundant than the rice fields in Location B. in the second planting period where rainfall had decreased quite sharply causing the condition of the paddy fields to become drier, but specifically at Location A the condition of the paddy fields during the second planting period remained at a planting interval so that it had higher humidity around the planting. This condition probably caused the predatory arthropod community to be more abundant than the rice fields in Location B. in the second planting period where rainfall had decreased quite sharply causing the condition of the paddy fields to become drier, but specifically at Location A the condition of the paddy fields during the second planting period remained at a planting interval so that it had higher humidity around the planting. This

condition probably caused the predatory arthropod community to be more abundant than the rice fields in Location B.

Table 1. Population characteristics of rice planthopper predatory arthropods by location and observation phase

No	Characteristics	Location A		Location B			
		MT I	Planting break	MT II	MT I	Planting break	MT II
1	Number of species	25	22	26	29	21	27
2	Total number of individuals	745	440	1.027	1,586	632	748
3	Species diversity index	2.09	2.38	2.40	2.02	2.14	2.33
4	Species evenness index	0.65	0.77	0.74	0.60	0.70	0.71

Information:

MT I : first planting season, 24 Sept – 31 Dec 2018(15 x observations)

Planting break: 7 Jan – 25 Feb 2019 (8 x observations)

MT II : second planting period, March 4 – June 17, 2019 (15 x observations)

The species diversity index has increased over time. The data in Table 1 shows the fact that the diversity of predatory arthropods in paddy fields is thought to be unaffected by the planting period because although the paddy fields experience a fairly long planting interval (two months), it does not decrease the species diversity index.

Figure 1 illustrates two interesting facts regarding the growing season. The first fact is an increase in the index over time without being influenced by the planting interval between the two planting periods, both at Location A and at Location B. This illustration shows that the planting interval does not cause a decrease in the diversity index of arthropod species in the paddy field ecosystem. Although the number of species during the planting interval decreased, it was offset by a decrease in the number of individuals so that the species diversity index showed an increase compared to the first planting period. Thus, it can also be stated that during the planting interval, the ecosystem conditions in paddy fields are maintained at a moderate level of balance.




Figure 1. Diversity index of predatory arthropod species by observation phase

The second fact is that the index at Location A is higher than that at Location B. The diversity of predatory arthropod species at Location A shows a higher index than Location B, whereas in terms of total individuals and species of predatory arthropods at Location B, it is higher than Location A. Species

diversity is a function of the species and the number of individuals per species. Theoretically, the abundance of certain species which is higher than other species will disproportionately lower the value of the species diversity index. This condition is in accordance with the observed facts.

The evenness index description is almost the same as the diversity index. Table 1. states that there is a tendency for the evenness index to increase from time to time without being influenced by the planting interval, both at Location A and at Location B. Even at Location B, evenness during the planting interval has the highest index compared to all planting phases. . Figure 2 can also illustrate that at each observation phase the evenness of species at Location A is higher than evenness at Location B.




Figure 2. Evenness index of predatory arthropod species by observation phase

Agroforestry has the potential to support the diversity of arthropod species in rice fields. The results of the analysis on the diversity of predatory arthropod species data based on observations during the two growing seasons (three phases of observation) illustrate that the agroforestry ecosystem around the rice fields is able to provide support for the diversity and even distribution of predatory arthropod species in the rice fields. Descriptive analysis of three observation phases (first planting period, planting interval and second planting period) showed that between the two planting periods interspersed with a fairly long planting interval did not reduce the diversity of predatory arthropod species in the rice field ecosystem. Vegetation structure in agroforestry, presence of grass in rice fields and in agroforestry,

Migration of several insect species from rice field ecosystems to agroforestry ecosystems occurs during the planting interval. The most preferred place for these arthropod species is the forest floor, which is mostly covered with grass, *Paspalum vaginalum* species, *Leersia* sp. and *Imperata cylindrica*. The existence of grass which is the choice of several rice pests as a migration destination will at the same time attract natural enemies to come to the same destination due to the availability of prey sources. Agroforestry in this case is an alternative habitat for arthropods that live in rice fields.

Agroforestry is a land management system that maintains three environmental elements. The social, economic and ecological elements in agroforestry land management have been unconscious for a long time, making this land management system superior and not found in other land management systems. Farmers get economic benefits in layers from the land they own. Agroforestry at the study site can be an example of this. Short-term needs are met from the forest by means of land planted with plant species of daily needs such as kencur, laja and cayenne pepper. Medium-term needs are represented by fruit species, namely cloves, rambutan, mango, salak and coconut. The long-term needs obtained by farmers from woody plants are teak, albasiah, mahogany or fruit plants that are quite old and unproductive to produce fruit. Farmers developing cropping patterns on their land

cannot be separated from the cultural elements inherent in their social environment so that between one area and another will have differences.

Local wisdom in farming communities was born from hereditary experiences. Cultivation carried out by farmers in West Java is also related to the local wisdom of the surrounding community. An example that can be observed is placing cattle pens around the forest so that there is a mutually beneficial relationship between the forest and livestock. Farmers get their source of animal feed from the forest environment and the forest benefits because fertilizer is available sustainably from the barn. Farmers also deliberately allow grass to grow on the forest floor because at certain times it becomes an easy and cheap source of animal feed. This condition will at the same time build a stronger ecosystem because the balance of ecological elements can be maintained by itself.

West Java farmers in many places place rice fields side by side with agroforestry. This situation is related to pragmatic reasons and environmental conditions itself. Farmers view that rice fields adjacent to an agroforestry environment can provide benefits, for example manure from livestock or forage from forests can be applied as fertilizer for rice plants in the fields. They can also do two main things efficiently at the same time, namely work in the garden and in the fields that are not far apart. The reason related to environmental conditions is because topographically the West Java region has a unique topography so that the villages are in a hilly condition.

The important value of this research is the exploration of ecological value and scientific intervention for practical value. This study tries to explore the conditions that develop in West Java farming communities where they place rice fields side by side with agroforestry. Many studies related to rice fields and agroforestry have been carried out, but very few studies have attempted to link the ecological value of forests to the interest of lowland rice cultivation. Farmers so far have been cultivating lowland rice side by side with agroforestry, not considering the scientific side in the form of ecosystem support for one another. The ecological point of view states that these two ecosystems allow for mutual support, although in a very simple context.

The scientific intervention for practical cultivation activities is to create an agroforestry area around the rice fields. The diversity of plants found in agroforestry is significant for the diversity of predatory arthropods in paddy fields. While theoretically, rice fields are an unequal ecosystem because the biotic and abiotic elements in it tend to be fragmented. Inequality of rice field ecosystem elements will be recovered by creating complementary ecosystems in the form of agroforestry in the vicinity. In this case, the agroforestry ecosystem becomes a complementary factor for the rice field ecosystem.

CONCLUSION

The study found that agroforestry was able to support the existence of predatory arthropods in rice fields. The existence of agroforestry around rice fields is able to provide support in the form of diversity of predatory arthropod species in a sustainable manner, both in simple agroforestry and in complex agroforestry. The diversity and evenness of arthropod species are not even affected by the season and planting intervals carried out by farmers in the fields. The results of this study, therefore, can reveal the ecological value of agroforestry for paddy fields, in relation to the sustainability of the diversity of predatory arthropod species in paddy fields.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to declare.

REFERENCES

1. Adnan, M. (2019). Keragaman Arthropoda Herbivora dan Musuh Alami Pada Tanaman Padi Lahan Rawa di Rowopulo Kecamatan Gumukmas Kabupaten Jember. *Jurnal Pengendalian Hayati*, 2(1), 10-16.
2. Aldini, G. M., Martono, E., & Trisyono, Y. A. (2019). Diversity of Natural Enemies Associated with Refuge Flowering Plants of *Zinnia elegans*, *Cosmos sulphureus*, and *Tagetes erecta* in Rice Ecosystem. *Jurnal Perlindungan Tanaman Indonesia*, 23(2), 285-291.
3. ALIFFAH, A. N., Natsir, N. A., RIJAL, M., & SAPUTRI, S. (2020). Pengaruh faktor lingkungan terhadap pola distribusi spasial dan temporal musuh alami di lahan pertanian. *Biosel: Biology Science and Education*, 8(2), 111-121.
4. Budiarti, L., Kartahadimaja, J., Sari, M. F., Ahyuni, D., & Ni Siluh Putu Nuryanti, D. (2021). Kelimpahan Relatif Artropoda Predator Penghuni Tajuk Tanaman Padi Galur Padi Rakitan Politeknik Negeri Lampung. *AgriHumanis: Journal of Agriculture and Human Resource Development Studies*, 2(1), 1-8.
5. Dadi, D. (2019). Local Ecology-Based Agroforestry Management: Building Effectiveness of Knowledge-Based Wetland Management. *Jurnal Mantik*, 3(3), 170-176.
6. Dadi, D. (2020). Community Characteristics of Brown Planthopper (*Nilaparvata lugens* Stal) Predator Arthropoda Community in Agroforestry-Based Rice Fields. *Budapest International Research in Exact Sciences (BirEx) Journal*, 2(4), 539-547.
7. Deosi, K. K., Suri, K. S., & Chandi, A. K. (2020). Effect of selection pressure of novel insecticides on brown planthopper, *Nilaparvata lugens* (Stål.) infesting rice. *Journal of Experimental Zoology, India*, 23(1), 71-75.
8. Desiska, S., Syahrawati, M., Arneti, A., Resti, Z., Martinius, M., Rahma, H., ... & Hidayat, R. (2019). Melatih Anggota Keltan Rambut dan Keltan Sakato Kota Padang untuk Mengendalikan Wereng Batang Coklat (WBC) dengan Joint Predator. *Jurnal Warta Pengabdian Andalas*, 26(4. a), 222-228.
9. Dhaliwal, S. S., Naresh, R. K., Walia, M. K., Gupta, R. K., Mandal, A., & Singh, R. (2019). Long-term effects of intensive rice-wheat and agroforestry based cropping systems on build-up of nutrients and budgets in alluvial soils of Punjab, India. *Archives of Agronomy and Soil Science*.
10. Dharshini, G. M., & Gowda, D. K. S. (2014). Biochemical basis of resistance in rice landraces to brown planthopper, *Nilaparvata lugens* (Stal.). *Current Biotica*, 8(3), 213-219.
11. Grimaud-Hervé, D., Balzeau, A., Widianto, H., Djubiantono, T., Détroit, F., Moigne, A. M., ... & Sémah, F. (2016). Position of the posterior skullcap fragment from Sendang Klampok (Sangiran Dome, Java, Indonesia) among the Javanese *Homo erectus* record. *Quaternary International*, 416, 193-209.
12. Hanif, K. I., Herlinda, S., Irsan, C., Karenina, T., Anggraini, E., Suwandi, S., & Susilawati, S. (2019).

- Populasi Serangga Hama dan Artropoda Predator pada Padi Rawa Lebak Sumatera Selatan yang Diaplikasikan Bioinsektisida dari *Beauveria bassiana* dan Insektisida Sintetik. *Jurnal Lahan Suboptimal: Journal of Suboptimal Lands*, 8(1), 31-38.
13. Isrin, M., & Fauzan, A. (2019). Pengaruh Frekuensi dan Saat Aplikasi *Beauveria bassiana* terhadap Wereng Batang Coklat (*Nilaparvata lugens* Stal) pada Tanaman Padi (*Oryza sativa* L.). *Biofarm: Jurnal Ilmiah Pertanian*, 14(2).
 14. Li, N., Chen, J., Zhang, J., He, Y., & Chen, L. (2011). Comparison for activities of antioxidant enzymes in imidacloprid resistant-and susceptible symbiotic strains in rice brown planthopper, *Nilaparvata lugens* (Stål). *Acta Phytopythologica Sinica*, 38(3), 258-264.
 15. Liu, X., & Lou, Y. (2018). Comparison of the defense responses in rice induced by brown planthopper *Nilaparvata lugens* (Stål) and white-backed planthopper *Sogatella furcifera* (Horváth). *Journal of Plant Protection*, 45(5), 971-978.
 16. Mahrub, E. (1998). Struktur komunitas artropoda pada ekosistem padi tanpa perlakuan pestisida. *Jurnal Perlindungan Tanaman Indonesia*, 4(1), 19-27.
 17. Meilin, A. (2017). Inventarisasi Parasitoid Telur Wereng Batang Cokelat di Beberapa Pertanaman Padi. *Jurnal Ilmiah Universitas Batanghari Jambi*, 12(3), 19-22.
 18. Muazam, A., & Nugroho, N. (2020). Efikasi Pestisida Hayati Pada Padi Varietas Tahan Tungro. *Jurnal Riset Agribisnis dan Peternakan*, 5(1), 30-38.
 19. Nasral, T. J., Syahrawati, M., & Liswami, Y. (2020). Daya Predasi dan Tanggap Fungsional Kumbang Unta (*Ophionea nigrofasciata*) pada Beberapa Kepadatan Wereng Batang Coklat (*Nilaparvata lugens*). *JPT: JURNAL PROTEKSI TANAMAN (JOURNAL OF PLANT PROTECTION)*, 4(1), 11-20.
 20. Ngatimin, S. N. A., Fatahuddin, F., Widarawati, R., & Nurfadila, N. (2020). Fluktuasi Populasi Wereng Coklat (*Nilaparvata lugens* Stal.) pada Tiga Macam Varietas Tanaman Padi (*Oryza sativa* L.). *BIOMA: JURNAL BIOLOGI MAKASSAR*, 5(2), 161-168.
 21. Nuryanti, N. S. P., Martono, E., & Ratna, E. S. (2018). The bioactivities of selected Piperaceae and Asteraceae plant extracts against brown plant hopper (*Nilaparvata lugens* Stål.). *Journal of ISSAAS (International Society for Southeast Asian Agricultural Sciences)*, 24(2), 70-78.
 22. Rasul, M. G., Widiyanto, W., & Mohanty, B. (2005). Assessment of the thermal performance and energy conservation opportunities of a cement industry in Indonesia. *Applied thermal engineering*, 25(17-18), 2950-2965.
 23. Rightmire, G. P., Margvelashvili, A., & Lordkipanidze, D. (2019). Variation among the Dmanisi hominins: multiple taxa or one species?. *American journal of physical anthropology*, 168(3), 481-495.
 24. RIZKIANI, S. H., & Herlinda, S. (2018). KEPEKAAN LARVA KUMBANG PREDATOR WERENG COKLAT (*Menochilus sexmaculatus*) TERHADAP *Cordyceps militaris*= THE SENSITIVITY OF BROWN PLANTHOPPER PREDATORY BEETLE (*Menochilus sexmaculatus*) TO *Cordyceps militaris* (Doctoral dissertation, Sriwijaya University).

25. Roy, S., Barooah, A. K., Ahmed, K. Z., Baruah, R. D., Prasad, A. K., & Mukhopadhyay, A. (2020). Impact of climate change on tea pest status in northeast India and effective plans for mitigation. *Acta Ecologica Sinica*, 40(6), 432-442.
26. Shepard, B. M. (1995). Rice-feeding insects of tropical Asia. *Int. Rice Res. Inst.*.
27. Silva, D. B., Jiménez, A., Urbaneja, A., Pérez-Hedo, M., & Bento, J. M. (2021). Changes in plant responses induced by an arthropod influence the colonization behavior of a subsequent herbivore. *Pest Management Science*.
28. Srinivasa, N., Chander, S., Sagar, D., & Venkatesh, Y. N. (2020). Rice brown planthopper prediction model with sweepnet catches. *Indian Journal of Entomology*, 82(3), 568-571.
29. Sugiyama, M., Yamamoto, E., Mukai, Y., Kaneko, Y., Nishizawa, M., & Harashima, S. (2006). Chromosome-shuffling technique for selected chromosomal segments in *Saccharomyces cerevisiae*. *Applied microbiology and biotechnology*, 72(5), 947-952.
30. Sun, Z., Yan, F., & Wang, M. (2018). Transgenic expression of Bt in rice does not affect feeding behavior and population density of the brown planthopper, *Nilaparvata lugens* Stål (Hemiptera: Delphacidae). *Entomologia Generalis*, 37(1), 35-45.
31. Supriyadi, S., & Wijayanti, R. (2018). Genetic similarity of brown planthopper, *Nilaparvata lugens* Stål. (Homoptera: Delphacidae) population from klaten and yogyakarta based on RAPD-PCR. *Indonesian Journal of Entomology*, 15(2), 85-92.
32. Tanaka, Y., Suetsugu, Y., Yamamoto, K., Noda, H., & Shinoda, T. (2014). Transcriptome analysis of neuropeptides and G-protein coupled receptors (GPCRs) for neuropeptides in the brown planthopper *Nilaparvata lugens*. *Peptides*, 53, 125-133.
33. Wang, J., Ding, B., Wang, L., Wang, Z., Zhou, G., & Lou, Y. (2018). The effects of exogenous castasterone treatment on rice brown planthopper *Nilaparvata lugens* behaviors, with special reference to feeding and ovipositing preferences. *Journal of Plant Protection*, 45(5), 998-1004.
34. Widiyanto, H., & Zeitoun, V. (2003). Morphological description, biometry and phylogenetic position of the skull of Ngawi 1 (East Java, Indonesia). *International Journal of Osteoarchaeology*, 13(6), 339-351.
35. Zeitoun, V., Barriel, V., & Widiyanto, H. (2016). Phylogenetic analysis of the calvaria of *Homo floresiensis*. *Comptes Rendus Palevol*, 15(5), 555-568.
36. Zeitoun, V., Déroit, F., Grimaud-Hervé, D., & Widiyanto, H. (2010). Solo man in question: Convergent views to split Indonesian *Homo erectus* in two categories. *Quaternary International*, 223, 281-292.

Predatory Arthropods For Brown Planthopper Species (Nilaparvata Lugens Stal): A Comparisional Study Based On Agroforestric

ORIGINALITY REPORT

5%

SIMILARITY INDEX

2%

INTERNET SOURCES

4%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Dewi Meidalima, Ruarita Ramadhalina Kawaty, Erlan B. Gunawan. "DIVERSITY OF ARTHROPOD PREDATOR IN SWAMP RICE FIELDS IN SOUTH SUMATERA", JURNAL HAMA DAN PENYAKIT TUMBUHAN TROPIKA, 2019
Publication 1%
- 2** Submitted to University of Basrah - College of Science
Student Paper 1%
- 3** rjoas.com
Internet Source 1%
- 4** Y Th Latupapua, F F Tetelay, F Soselisa, C K Pattinasarany, L Latupapua, A Sanduan. "In-situ kuskus conservation strategy in Wakasihu Village, West Leihitu District, Central Maluku District", IOP Conference Series: Earth and Environmental Science, 2021
Publication <1%

5

S K Aminuddin, B Nurkin, S Millang, S H Larekeng. "Function of tree components in agroforestry forms in Bolaromang Village, Buttono District, Gowa Regency", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

6

journal.ugm.ac.id

Internet Source

<1 %

7

Ligar Novi Ayuniar, Jafron Wasiq Hidayat. "The Profile Quality of Pond In Kendal Regency to Diversification Aquaculture", E3S Web of Conferences, 2018

Publication

<1 %

8

Wahyu Prihanta, Facischa Ayu Irviandari, Elly Purwanti, Abdulkadir Rahardjanto, Fuad Jaya Miharja. "Contamination of shrimp pond waste: The impact on macrozoobenthos diversity", AIP Publishing, 2020

Publication

<1 %

9

analesdepediatria.org

Internet Source

<1 %

10

journals.plos.org

Internet Source

<1 %

11

www.frontiersin.org

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On