

Kurniati Devi Purnamasari

ARTIKEL JMRH

 MANUSKRIP
 devi
 Universitas Galuh

Document Details

Submission ID**trn:oid::1:3641638626****Submission Date****Sep 6, 2026, 3:50 PM UTC****Download Date****Sep 6, 2026, 3:52 PM UTC****File Name****Artikel_JMRH_Kurniati_Devi_P.pdf****File Size****468.7 KB****21 Pages****7,038 Words****39,512 Characters**

27% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Filtered from the Report

- Bibliography
- Quoted Text

Match Groups

-  **109 Not Cited or Quoted 22%**
Matches with neither in-text citation nor quotation marks
-  **28 Missing Quotations 5%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 22%  Internet sources
- 20%  Publications
- 5%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- 109 Not Cited or Quoted 22%**
Matches with neither in-text citation nor quotation marks
- 28 Missing Quotations 5%**
Matches that are still very similar to source material
- 0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 22% Internet sources
- 20% Publications
- 5% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Internet	
ijnhs.net		2%
2	Internet	
perpustakaan.poltekkes-malang.ac.id		1%
3	Publication	
Marta Armita Br Silaban, Novita Andriani Br Manjorang, Debby Chintya Yun, Erin ...		1%
4	Internet	
www.frontiersin.org		<1%
5	Publication	
Risa Pitriani, Miratu Megasari. "Effectiveness of Oxytocin Massage on Increasing ...		<1%
6	Internet	
repository.ar-rum.ac.id		<1%
7	Internet	
docksci.com		<1%
8	Internet	
www.sciencegate.app		<1%
9	Publication	
Rahmadini Faricha Hakim, Dyan Permata Rachim, Hindun Mila Hudzaifah, Ulfa Fa...		<1%
10	Internet	
jmrh.mums.ac.ir		<1%

11	Internet	www.researchsquare.com	<1%
12	Student papers	FACULTY OF SOCIAL SCIENCES (FSS)	<1%
13	Internet	pmc.ncbi.nlm.nih.gov	<1%
14	Internet	academic.oupdev.silverchair.com	<1%
15	Internet	journal.uin-alauddin.ac.id	<1%
16	Internet	ejournal2.unud.ac.id	<1%
17	Internet	www.ncbi.nlm.nih.gov	<1%
18	Publication	Cemile Kütmeç Yılmaz, Güler Duru Aşiret, Funda Çetinkaya. "The effect of back m...	<1%
19	Publication	Elif Dağlı, Neşe Çelik. "The effect of oxytocin massage and music on breast milk p...	<1%
20	Publication	G Attilakos. "Carbetocin versus oxytocin for the prevention of postpartum haemo...	<1%
21	Publication	Pebia Norviana, Sunartono Sunartono, Hargianti Dini Iswandari, Siti Maimunah, I...	<1%
22	Internet	nutrition.bmj.com	<1%
23	Internet	sci-hub.red	<1%
24	Internet	tpmap.org	<1%

25	Publication	Morten Møller. "Vasopressin and oxytocin beyond the pituitary in the human brai...	<1%
26	Student papers	Saveetha Dental College and Hospital, Chennai	<1%
27	Internet	kb.psu.ac.th:8080	<1%
28	Internet	oamjms.eu	<1%
29	Internet	www.e-mjm.org	<1%
30	Internet	www.rba.periodikos.com.br	<1%
31	Publication	Luigi Taranto-Montemurro, Bradley A. Edwards, Scott A. Sands, Melania Marques,...	<1%
32	Student papers	University of Wolverhampton	<1%
33	Internet	dataprojectng.com	<1%
34	Publication	Ayşenur Sariaslan, Mehtap Kavurmaci. "Evaluation of the effect of the PRECEDE P...	<1%
35	Publication	Silvia Mona, Kiki Rizki Dasaryandi, Susanti Susanti. "The Effect of Combined SEFT ...	<1%
36	Internet	assets-eu.researchsquare.com	<1%
37	Internet	jurnal.ugm.ac.id	<1%
38	Internet	www.eelet.org.uk	<1%

39	Student papers	Universitas Galuh	<1%
40	Internet	ijahst.org	<1%
41	Publication	Anna Coutsoudis, Penny Reimers, John Cassey, Tanya Doherty. "Wet Nursing and ...	<1%
42	Publication	Kadek Yuli Hesti, Noor Pramono, Sri Wahyuni, Melyana Nurul Widyawati, Bedjo S...	<1%
43	Student papers	The University of Texas at Arlington	<1%
44	Internet	erickimphotography.com	<1%
45	Internet	healthdocbox.com	<1%
46	Internet	ijponline.biomedcentral.com	<1%
47	Internet	journals.kmanpub.com	<1%
48	Internet	sites.kowsarpub.com	<1%
49	Publication	Aya Asfandiyar Muhammed Saeed, Halgord Ali M. Farag, Nabil Hussain Rasul. "Im...	<1%
50	Internet	core.ac.uk	<1%
51	Internet	files01.core.ac.uk	<1%
52	Internet	public-pages-files-2025.frontiersin.org	<1%

53	Internet	www.mdpi.com	<1%
54	Publication	Astrida Budiarti, Miftaql Jannah, Puji Hastuti. "A Pre Experimental Study: The Eff...	<1%
55	Publication	Fateme Norouzi Nodehi, Parvin Aziznejadroshan, Hossein-Ali Nikbakht, Mohsen H...	<1%
56	Publication	Mo Chen, Yaqiong Cui. "The effects of AWE and peer feedback on cohesion and co...	<1%
57	Publication	Rahmi Amtha, Ferry Sandra, Rosalina Tjandrawinata, Indrayadi Gunardi, Anggrae...	<1%
58	Internet	narraj.org	<1%
59	Publication	Kurnia Dewiani, Riana Versita , Fitri Ramadhaniati , Endah Purda Listya . "The Ef...	<1%
60	Publication	Sumarni Sumarni, Yuni Purwati. "A case study of oxytocin massage intervention f...	<1%
61	Internet	bmcpregnancychildbirth.biomedcentral.com	<1%
62	Internet	dehesa.unex.es	<1%
63	Internet	digitalcommons.uconn.edu	<1%
64	Internet	jpcp.uswr.ac.ir	<1%
65	Internet	mdpi-res.com	<1%
66	Internet	repository.pkr.ac.id	<1%

67	Internet	research.library.mun.ca	<1%
68	Internet	www.iris.unict.it	<1%
69	Publication	Şeyma Kılçı Erciyes, Oya Kavlak. "The effect of back and breast massage on the a...	<1%
70	Publication	Adinda Putri Sari Dewi, Kusumastuti Kusumastuti. "Effect of combining oketani a...	<1%
71	Publication	Kaja Looock, Lars Schwabe. "Cognitive training prevents stress-induced working m...	<1%
72	Publication	Lori Feldman-Winter, Sahira Long, Julie Ware, Margaret G. Parker et al. "Opportu...	<1%
73	Publication	Savas Karakus, Halef Okan Dogan. "Exploring altered free amino acids and metab...	<1%
74	Publication	Sulastry Pakpahan, Hetty Panggabean. "Herbal Tea of Torbangun Leaf and Exclus...	<1%
75	Publication	Tri Ratna Ariestini, Christine Aden, Berthiana Berthiana, Harlyanti Muthma'innah...	<1%
76	Publication	Xiangyu Tan, Peiyang Mao, Hongfan Yu, Li Tang et al. "Translation and psychome...	<1%
77	Internet	advbiores.mui.ac.ir	<1%
78	Internet	cdn.techscience.cn	<1%
79	Internet	journal.universitaspahlawan.ac.id	<1%
80	Internet	www.mltj.online	<1%

JMRH-2412-2550

The Effect of 30-Minute Oxytocin Massage on Breast Milk Volume and Infant Weight Gain in Early Postpartum Women: A Randomized Controlled Trial

Kurniati Devi Purnamasari^{1*}, Yudita Ingga Hindiarti¹, Erna Yovi Kurniawati², Iqbal Pramukti³

¹Department of Midwifery, Faculty of Health Sciences, Universitas Galuh, Indonesia

²Department Midwifery Profession, Alma Ata University, Yogyakarta, Indonesia

³Department of Community, Faculty of Nursing, Universitas Padjajaran, Indonesia

Running title : *Oxytocin Massage on Milk Production and Infant Growth*

Abstract

Background and aim: Many new mothers discontinue breastfeeding prematurely. Exclusive Breastfeeding (EBF) during the first six months of life is widely recognized as an optimal feeding practice that supports infant health and survival. This study aimed to determine the effect of a 30-minute oxytocin massage on breast milk volume and infant weight gain among early postpartum women.

Methods: A randomized single-blind controlled trial with a two-group pretest–posttest design was conducted among 60 postpartum women. Participants were randomly assigned using computer-generated randomization with allocation concealment. The intervention group received a 30-minute oxytocin massage, while the control group received a 15-minute oxytocin massage for three consecutive days. Breast milk volume was measured using standardized manual milk expression, and infant weight was assessed using calibrated infant weighing scales. Outcomes were evaluated at baseline and after completion of the intervention. Data were analyzed using paired- and independent-samples *t*-tests at $\alpha = 0.05$.

Results: The mean age of participants was 28.3 ± 5.1 years. The increase in breast milk volume and infant weight was greater in the intervention group (mean = 10.03 mL and 151.50 g) than in the control group (mean = 8.33 mL and 106.16 g). These differences were statistically significant

for both breast milk volume ($t = -8.171, p < .001$) and infant weight gain ($t = -9.088, p < .001$). ANCOVA also showed significant between-group differences.

Conclusion: A 30-minute oxytocin massage may improve breast milk volume and infant weight gain, offering a practical, low-cost method for postpartum lactation support.

Keywords: Postpartum Period; Breastfeeding; Lactation; Massage; Infant Growth

Introduction

Breastfeeding is a crucial intervention for reducing infant mortality and promoting optimal child health (1). In impoverished areas, providing exclusive breast milk saves approximately 1.5 million newborns annually (2). Globally, only 48% of infants under six months of age are exclusively breastfed, remaining below the World Health Assembly target of 60% by 2030 (2) (3). Acknowledging its critical role, the World Health Organization endorses exclusive breastfeeding for the first six months as a means of supporting infant growth, development, and overall health (4,5).

Data from the WHO in 2021 indicated that the global rate of exclusive breastfeeding for infants was only around 44 percent (5). Furthermore, only 67.74% of 0 to 6-month-old Indonesian babies received exclusively breast milk, compared to the target of 80%. Exclusive breastfeeding was reported in only 63.35% of infants based on data from the Health Office of West Java Province (6). Ciamis District achieved the second lowest score at 45.96%, still below the target (6). In addition, reports from the Health Office of Ciamis District showed that the achievement of exclusive breastfeeding in 2019 did not meet the intended target, with coverage limited to 49% (7). This situation was especially prevalent in the working area of Baregbeg Public Health Center, where exclusive breastfeeding rates remain below the national target. This problem results from factors such as a lack of awareness and insufficient healthcare support.

Assessing the efficacy of oxytocin massage in this specific circumstance is essential, considering that the procedures and obstacles differ in Indonesia (8). Despite these benefits, exclusive breastfeeding (EBF) faces several challenges, including maternal concerns about insufficient

breast milk production (9). Many mothers experience difficulties in maintaining an adequate milk supply (10), which can lead to early supplementation with formula or other substitutes. Addressing these challenges is essential to improving breastfeeding rates and reducing infant mortality (11).

Breast milk production is regulated by hormonal mechanisms, particularly the role of oxytocin (12). Oxytocin massage is a proven method to enhance milk production by stimulating the spinal cord, activating neurotransmitters in the medulla oblongata, and signalling the hypothalamus (13). This process triggers the posterior pituitary gland to release oxytocin, which facilitates milk ejection and sustains lactation (14). Breastfeeding provides nutritional and emotional benefits for both children and mothers (15).

Furthermore, long-term benefits include a lower risk of breast and uterine cancers, obesity, osteoporosis, and type 2 diabetes mellitus (16,17). However, breastfeeding can be more challenging since it depends on the production of breast milk (18). Breastfeeding is facilitated by adequate breast milk production (18). Inadequate breast milk for the baby can result from a lack of or late production of breast milk (19). The two elements that affect breast milk production are production and release(20). The prolactin hormone influences breastmilk production; however, prolactin itself does not directly relate to the level of milk production (21). It is the oxytocin hormone that increases the movement of breastmilk to the breasts (22).

Existing solutions to address breastfeeding challenges include pharmacological interventions, breastfeeding support programs, and maternal education (23). Pharmacological treatments are occasionally recommended to supplement lactation despite their potential side effects for both women and babies (24). Breastfeeding support, including professional lactation counseling and peer support groups, has been proven effective in improving breastfeeding outcomes by addressing common concerns and enhancing maternal confidence (25). Additionally, educational programs empower mothers with adequate knowledge and skills about breastfeeding and strategies to overcome common lactation difficulties (26).

Oxytocin massage is a non-pharmacological intervention designed to stimulate the secretion of oxytocin (27). Oxytocin massage and proper breast care may augment breast milk production, with efficacy contingent upon the frequency and duration of breastfeeding (18). This intervention

21 activates the parasympathetic nervous system, thereby promoting oxytocin release from the posterior pituitary, particularly when massage is applied along the spinal area from the fifth to sixth ribs toward the scapular region (28).

9
3 However, non-pharmacological methods like oxytocin massage offer a safer, more natural, and cost-effective approach to supporting lactation (27,29). Oxytocin massage stimulates the release of oxytocin, which enhances milk ejection and overall milk production without the risks associated with medication (30). Stimulation of the parasympathetic nervous system through oxytocin massage and spinal massage along the fifth to sixth ribs extending to the scapular region accelerates oxytocin release from the posterior pituitary (28). Besides being easy to implement, either by trained healthcare providers or mothers, this technique has been shown to improve breastfeeding success rates (31). Given its accessibility, effectiveness, and minimal risk, oxytocin massage presents an innovative solution to breastfeeding challenges, particularly in regions with limited healthcare resources.

42
6
8 Data from the WHO in 2021 indicated that the global rate of exclusive breastfeeding for infants was only around 44 percent (World Health Organization, 2021). Furthermore, only 67.74% of 0 to 6-month-old Indonesian babies received exclusively breast milk, compared to the target of 80%. Exclusive breastfeeding was reported in only 63.35% of infants based on data from the Health Office of West Java Province (6). Ciamis District achieved the second lowest score at 45.96%, still below the target (6). In addition, reports from the Health Office of Ciamis District showed that the achievement of exclusive breastfeeding in 2019 did not meet the intended target, with coverage limited to 49% (7).

6
4 5 This situation was especially prevalent in the working area of Baregbeg Public Health Center, where exclusive breastfeeding rates remain below the national target. This problem results from factors such as a lack of awareness and insufficient healthcare support. This study aimed to determine the effect of a 30-minute oxytocin massage on both breast milk volume and infant weight gain in early postpartum women.

Materials and Methods

43 This study employed a randomized single-blind controlled trial with a two-group pretest–posttest design involving 60 postpartum women at 14 days postpartum at Baregbeg Public Health Center,

Ciamis, West Java, Indonesia. Data collection was conducted from July to November 2020. Eligible participants were postpartum mothers who were breastfeeding their infants at 14 days after vaginal delivery, had assumed care of their infants after hospital discharge, were able to provide informed consent, were not taking any medications, and had no history of spinal injury. Women were excluded if they had a history of cesarean delivery, were unable to communicate in Indonesian, or their infants required special care for more than 24 hours. The intervention was administered at home for three consecutive days, from postpartum day 14 to day 16. Baseline measurements were obtained before the first massage session on postpartum day 14, and post-intervention measurements were obtained after completion of the third massage session on postpartum day 16.

Sample size estimation was conducted using G-power analysis. The calculation was based on the t-test family, assuming a large effect size (0.8), an alpha error probability of 0.10, and a statistical power of 0.90. The minimum sample size was 56 subjects. The assumption of a large effect size was based on published methodological literature that outlines acceptable strategies for effect size selection in quantitative studies, particularly in cases where prior empirical estimates or pilot data are not available (32). This effect size was selected to ensure robust results and to reflect the potential impact of the 30-minute oxytocin massage on breast milk volume and infant weight gain. The minimum target sample size was 56 participants, with an alpha error probability of 0.1 and a power of 0.9, as calculated using G*Power analysis. This approach ensures the credibility and reliability of the study's findings.

The study used a two-group randomized single-blind controlled trial. The random sequence was generated using a computer-based randomization procedure by a trained enumerator who was not involved in participant eligibility assessment or recruitment. Allocation concealment was ensured using sequentially numbered, opaque, sealed envelopes prepared by the same independent enumerator. Group assignments were revealed only after participant enrollment had been completed, thereby minimizing selection bias. Outcome assessors were blinded to group allocation to reduce potential assessment bias. A total of 30 participants were allocated to the control group and received a 15-minute oxytocin massage, while 30 participants were allocated to the intervention group and received a 30-minute oxytocin massage. Both interventions followed the

same standardized oxytocin massage protocol, with the duration of massage being the only difference between the two groups.

71 A total of 60 participants provided written informed consent before study enrollment and received compensation for their participation. Each session used a moderate-pressure back massage. The intervention was delivered by the same eight massage therapists to standardize the type of touch among participants in each group. Screening methods, including questionnaires, were implemented to reduce the risk of disease transmission, and precautions were in place to ensure the safety of populations in affected communities during data collection. In addition, measures such as wearing personal protective equipment (PPE) and disinfecting equipment were applied to reduce the risks of disease transmission. A 30-minute massage or pressure session was administered to this group, focusing on the back of the neck, spinal area, and the fifth and sixth costal regions near the scapula with circular motions. Massage took place in their home, and the subjects attended daily sessions on days 1 to 3.

Eight massage therapists were involved in delivering the intervention. Before data collection, all therapists received standardized training from the research team on the oxytocin massage procedure, including participant positioning, massage sequence, massage technique, duration, and application of baby oil. The training included a demonstration and supervised practice to ensure that all therapists followed the same standardized protocol. The research team provided a written protocol and checklist to guide the implementation of each massage session. Adherence to the standardized massage protocol was monitored using the checklist during each session, covering participant positioning, massage sequence and technique, massage duration, and application of baby oil. The research team also used tracking sheets to monitor the completion of daily intervention sessions and maintained consistency in protocol implementation throughout the study. Therapists were instructed to minimize communication with participants during the intervention, and the same therapists were involved in each session to maintain consistency. Participants were informed beforehand that verbal interaction would be restricted during the intervention. The research team also used tracking sheets to monitor daily data collection during the pretest and posttest. No adverse effects associated with the massage intervention were reported in either group.

A personal information form was employed to acquire the data and questionnaires at the patients' homes, specifically in their rooms. The entire data collection process lasted about an hour. A personal information form was completed by the researcher. The instrument comprised 12 items assessing participants' medical and sociodemographic characteristics. The intervention began after the subjects were randomized.

A total of 60 women who met the study eligibility criteria completed the questionnaire and observation sheet administered by the interviewer. A structured questionnaire was developed based on a review of the relevant literature and expert consultation to collect participants' sociodemographic and obstetric characteristics. Twenty women who were excluded from the main study participated in a pilot test to assess the clarity and feasibility of the questionnaire. The first section collected information on participants' age, education, delivery characteristics, occupation, body mass index (BMI), type of residence, parity, and breast care. The second section consisted of an observation sheet completed by the enumerator to record infant weight before and after the intervention and breast milk volume measurements.

Both groups of patients received daily oxytocin massages for 15 minutes and 30 minutes, respectively, between 07:00 and 10:00 at home for three days. Before the massage, the patients and their environment were prepared, and privacy was ensured. The most comfortable position for the back massage was established through discussion with the participants. Each participant was seated on a chair with a side table positioned in front, and pillows were placed on the table to provide adequate support. Before initiating the massage, the researcher, who was trained in massage techniques, washed, dried, and warmed her hands, then applied 5–10 drops of room-temperature liquid baby oil, commonly used to moisturize the back, to her hands. The massage procedure was subsequently initiated, with the oil serving to lubricate the skin and minimize friction.

Breast milk volume was assessed by collecting the breast milk released during the standardized observation period into a calibrated glass container. The volume collected was recorded in milliliters (mL). The measurement was performed 2 hours after the participant's last breastfeeding session. This measurement represented the amount of breast milk released during the specified observation period and was not intended to represent total breast milk production over the entire

lactation period. The same observation period, calibrated container, and measurement procedures were used for all participants to ensure consistency.

Baseline measurements of breast milk volume and infant weight were obtained before the first massage session on postpartum day 14. The intervention was administered for three consecutive days, from postpartum day 14 to day 16. Follow-up measurements of breast milk volume and infant weight were conducted after completion of the third massage session on postpartum day 16 using the same measurement procedures and equipment. Thus, the follow-up period covered three days from the first to the final intervention session.

The research was approved by Dian Nuswantoro University HREC under protocol approval 319/EA/KEPK-Fkes-UDINUS/XI/2022. The data were collected at the Baregbeg Public Health Center, Ciamis, West Java, Indonesia. All subjects provided written informed consent after the procedures were explained. A total of 60 subjects participated in the study. This was a randomized single-blind clinical trial. It was registered with the code INA-100220250161FA1 in the Indonesia Clinical Research Registry (INA-CRR) by the Health Policy Agency of the Ministry of Health of the Republic of Indonesia.

A descriptive analysis was used to summarize participant characteristics. Data normality was assessed using the Kolmogorov–Smirnov test. The independent and dependent sample T-tests were conducted to compare the means of breast milk production before and after treatment, as well as between the groups. SPSS software version 20 for Windows was used. The exclusion criteria were checked during the study, and participants who met these criteria were excluded.

The research was carried out at the Public Health Center in Ciamis City, Indonesia.

Sociodemographic and obstetric data were collected using a structure questionnaire developed based on relevant literature and expert consultation. The questionnaire included information on age, education, occupation, body mass index (BMI), parity, place of residence, and birth-related characteristics. Data collection was conducted through interviewer-administered questionnaires, direct observation, breast milk volume measurement, and infant weight assessment (33)

Result

The randomization flowchart is demonstrated in Figure 1. Approximately 59% of invited women were eventually randomised in the study; 31% of the invited women were less than 2 weeks postpartum, and 4% had a caesarean section. A total of 60 subjects were randomized in the study and analyzed.




Figure 1. CONSORT flow diagram of participants' selection

Overall, the 2-week postpartum women recruited were primarily adults with a mean age of 28.3 years (± 5.1) (Table 1). In terms of educational attainment, most participants had completed high school (40%) or junior high school (38.3%). The vast majority of participants (91.7%) had given early breastfeeding. For the parity status, most subjects were at parity 2. The body mass index (BMI) of the subjects was mainly in the healthy weight and overweight ranges (53.3% vs. 40%, respectively). Subjects living in the rural area were greater than those in the suburban area (80% vs. 20%, respectively). Compared to working mothers, non-working mothers were greater (88.3 vs. 11.7%, respectively). The mean age of the infants was 14.1 days (± 1.3).

For the intervention group, the mean maternal age was 28.8 (± 4.8), whereas the control group had a mean age of 27.8 ± 5.4 years 27.8 (± 5.4) (Table 1). Regarding educational attainment, most participants had completed senior high school (21.7%) and junior high school (21.7%).

Early initiation of breastfeeding was reported by nearly all participants (90%). In terms of parity, parity two was the most common status among participants. The body mass index (BMI) of the subjects was mainly in overweight and healthy weight status (53.3% vs 43.4%, respectively). Subjects living in the rural area were greater than those in the suburban area (90 45% vs. 10%, respectively). Compared to working mothers, non-working mothers were greater (86.7 vs. 13.3%, respectively). The mean age of the infants was 14.3 days (± 1.4).

Table 1. Socio-demographic characteristics of participants in the intervention and control groups

Characteristics	total (n= 60)	Control Group (n = 30) n (%)	Intervention Group (n = 30) n (%)	p value
Age (years) (mean $\pm$ SD)	28.3 $\pm$ 5.1	27. 8 $\pm$ 5.4	28.8 $\pm$ 4.8	.014
<=24 years	15 (25)	10 (16.6)	5 (8.4)	
>24 years	45 (75)	20 (33.3)	25 (41.7)	
Educational level				.248
Elementary	7 (11.7)	6 (10)	1 (1.7)	
Junior High School	23 (38.3)	10 (16.6)	13 (21.7)	
Senior High School	24 (40)	11 (18.3)	13 (21.7)	
University	6 (10)	3 (5)	3 (5)	
Birth-related characteristic				.014

Characteristics	total (n= 60)	Control Group (n = 30) n (%)	Intervention Group (n = 30) n (%)	p value
Vaginal delivery	60 (100)	36 (60)	24 (40)	
Preterm delivery	2 (3.4)	1 (1.7)	1 (1.7)	
Breast feeding	60 (100)	30 (50)	30 (50)	.542
Early breastfeeding	55 (91.7)	28 (46.7)	27 (45)	
Non-early breastfeeding	5 (8.4)	2 (3.4)	3 (5)	
Therapy				.282
Breast care	10 (16.7)	6 (10)	4 (6.7)	
Non-breast care	50 (83.3)	24 (40)	26 (43.3)	
Parity				.648
1 st -2 nd	49 (81.6)	25 (41.6)	24 (40)	
3 rd -5 th	11 (18.4)	5 (8.4)	6 (10)	
BMI				0.421
Normal Weight	33 (55)	19 (31.6)	14 (23.4)	
Overweight	27 (45)	11 (18.3)	16 (26.7)	
Type of residence				<.001
Rural (n)	48 (80)	21 (35)	27 (45)	
Suburban (n)	12 (20)	9 (15)	3 (5)	
Occupation				.591
Not working	53 (88.3)	27 (45)	26 (43.3)	
Working	7 (11.7)	3 (5)	4 (6.7)	
Age of infant (days) (mean ±SD)	14 ± 1.4	14.1 ± 1.3	14.3 ± 1.4	.620

Data were described as a mean ± SD and number (percent), P-value was calculated using the Chi-square test.

After receiving the standard protocol (15 minutes of oxytocin massage), subjects' breast milk volume increased significantly from 7.59 ml to 15.92 ml with a t-value of -10.91. For the infant's weight, there was also a significant improvement from 3571.67 grams to 3677.83 grams with a t-value of 5.68.

As shown in Table 2, the control group demonstrated a significant increase in breast milk volume, from 7.59 ± 3.05 mL before the intervention to 15.92 ± 5.08 mL after the intervention ($t = -10.913$, $p < .001$). Infant weight also increased significantly from 3571.67 ± 573.50 g to 3677.83 ± 579.69 g ($t = -5.683$, $p < .001$).

Table 2. The Effect of 15-Minute Oxytocin Massage on Breast Milk Volume among Postpartum Women (Control Group) (n=30)

Variable	Pre-intervention		Post-intervention		t*	p value
	Mean	SD	Mean	SD		
Breast milk(ml)	7.59	3.05	15.92	5.08	-10.913	< .001
Infant's weight(gr)	3571.67	573.50	3677.83	579.69	-5.683	< .001

*Statistical analysis was performed using a paired t-test.

A paired *t*-test analysis demonstrated that a 30-minute oxytocin massage significantly increased breast milk volume among postpartum women in the intervention group (Table 3). The mean breast milk volume increased from 8.75 ± 3.29 mL before the intervention to 18.78 ± 7.81 mL after the intervention ($t = -8.171, p < .001$). Similarly, the infants' mean body weight increased significantly from 3722 ± 544.57 grams to 3923.50 ± 91.30 grams following the intervention ($t = -9.088, p < .001$). These findings indicate that oxytocin massage was effective in enhancing both maternal milk volume and infant weight gain.

Table 3. The Effect of 30-Minute Oxytocin Massage on Breast Milk Volume among Postpartum Women (Intervention Group) (n=30)

Variable	Pre-intervention		Post-intervention		t*	p value
	Mean	SD	Mean	SD		
Breast milk (ml)	8.75	3.29	18.78	7.81	-8.171	<.001
Infant's weight (gr)	3722	544.57	3923.50	91.30	-9.088	<.001

*Statistical analysis was performed using a paired t-test.

Among the two groups, the independent *t*-test results showed a significant mean difference in breast milk after receiving the intervention.

Table 4 demonstrates a comparison of breast milk volume and infant weight between the intervention and control groups. The results showed that, with a mean difference of 1.70, the intervention group produced more breast milk on average ($M = 10.03, SD = 6.72$) than the control group ($M = 8.33, SD = 4.11$). In a similar vein, the mean weight difference between the infants in the intervention group ($M = 151.50, SD = 91.30$) and the control group ($M = 106.16, SD = 102.31$)

was 45.34. Partial eta squared ($\eta^2 = .423$) indicated a moderate effect size, and both outcomes showed statistically significant differences ($p < .001$). These findings imply that the intervention significantly increased infant weight and breast milk volume.

Table 4. The Comparison of breast milk volume and Infant's Weight between the Intervention and Control Groups (n=60)

Variable	Control Group		Intervention Group		Mean difference	Partial Eta Squared	p value
	Mean	SD	Mean	SD			
Breast milk (ml)	8.33	4.11	10.03	6.72	1.70	.423	< .001
Infant's weight (gr)	106.16	102.31	151.50	91.30	45.34	.423	< .001

Data are presented as mean and standard deviation (SD). Differences between groups were analyzed using an independent samples t-test.

Based on Table 5, the univariate ANCOVA indicated that, after controlling for pretest breast milk volume scores, there was a statistically significant effect of group on posttest breast milk volume ($p < .001$), showing a large effect size. Pretest scores were also a significant covariate ($p = .016$), meaning baseline breast milk volume scores significantly predicted post-intervention outcomes. These results indicated that the improvement in breast milk volume (710.12 ml, $p < 0.001$) had a strong and significant effect on the outcome, even after adjusting for baseline differences.

Table 5: ANCOVA for Post-Intervention breast milk volume Controlling for Baseline Scores

	Sum of Squares	df	Mean Square	F	p	Partial η^2
Pre-test (Covariate)	105.875	1	105.875	7.213	.016	0.311
Group	710.120	1	710.120	48.387	<.001	0.751
Error	235.446	16	14.676			
Total	1051.441	18				

According to the findings presented in Table 6, an ANCOVA result was carried out to examine differences in infants' posttest weight between the intervention and control groups, adjusting for pretest weight.

The results indicated that baseline weight was a significant predictor of post-intervention weight, $F(1, 16) = 8.35$, $p = .012$, partial $\eta^2 = .343$, suggesting a moderate contribution of baseline weight to the outcome. After controlling for this covariate, a significant difference emerged between the two groups, $F(1, 16) = 37.84$, $p < .001$, partial $\eta^2 = .703$. These findings indicate that the 30-minute oxytocin massage intervention was associated with significantly greater infant weight outcomes compared with the control group.

Table 6. ANCOVA linearity test of pre-test and post-test scores of the infant's weight

	Sum of Squares	df	Mean Square	F	p	Partial η^2
Pre-test (Covariate)	540.160	1	540.160	8.345	.012	0.343
Group (Control vs Intervention)	2450.560	1	2450.560	37.840	<.001	0.703
Error	1030.000	16	64.375			
Total	3471.220	19				

Discussion

This study aimed to evaluate the effect of a 30-minute oxytocin massage on breast milk volume and infant weight gain among early postpartum women. The findings of this study indicate that breast milk volume increased following three days of 30-minute oxytocin massage. These results are consistent with previous research demonstrating the beneficial effect of this intervention on enhancing breast milk production (34). However, the current study showed a higher mean difference in breast milk volume before and after receiving the intervention (10.03 vs. 8.33). Another study showed that oxytocin massage has many benefits in the breastfeeding process (35), such as increasing the milk production of postpartum mothers who have less milk production (36), reducing stress on postpartum mothers, reducing spinal pain, stimulating the work of the oxytocin hormone, enhancing comfort, improving the movement of milk to the breasts, increasing the frequency of breast filling, and supporting breastfeeding.

Massage was associated with elevated oxytocin levels and reductions in adrenocorticotropin hormone (ACTH), nitric oxide (NO), and beta-endorphin (BE). When compared with the rest

group, the massage group demonstrated significantly greater changes in oxytocin, ACTH, NO, and BE (37). Oxytocin is synthesized in the supraoptic and paraventricular nuclei of the hypothalamus and subsequently transported to the pituitary gland via the tuber cinereum and infundibulum (38) during oxytocin massage for approximately 30 minutes.

Oxytocin massage is a treatment associated with breast milk production. The intervention group showed a significantly greater increase in breast milk volume than the control group, suggesting that a longer duration of oxytocin massage may enhance lactation outcomes. This is in line with a study by Ike Johan and Ninik Azizah, who found higher odds of breast milk among mothers who received oxytocin massage compared to those in the control group who did not receive oxytocin massage (28). This is similar to the study findings from the theory that explains that giving a massaging the whole spine (vertebrae) stimulates the posterior pituitary for the secretion of oxytocin (39,40).

An intervention group given a 30-minute oxytocin massage showed a significant increase in the infant's weight gain, which was 1.42 times more likely to increase than the control group. This is in line with another study that found an increase in infants' weight gain in the first six months of their lives (41,42).

This study confirms that a 30-minute oxytocin massage significantly increases breast milk volume compared to standard care, highlighting its potential as a non-pharmacological intervention to support lactation. Given its simplicity, cost-effectiveness, and ease of implementation, it is recommended that oxytocin massage be integrated into routine postpartum care, particularly in low-resource areas with limited access to lactation support. Healthcare providers, including midwives and lactation consultants, could be trained to administer this technique and educate mothers on how to do it to optimize breastfeeding outcomes. This is in line with another study that found an increase in infants' weight gain in the first six months of their lives (41,42).

The findings of this study, supported by multiple prior studies, consistently underscore the may improve of oxytocin massage in enhancing breast milk volume and promoting infant weight gain. This intervention not only stimulates the physiological release of oxytocin but also contributes to maternal comfort, stress reduction, and improved lactation outcomes (43,44). These cumulative

findings reinforce the value of oxytocin massage as a practical, low-cost, and evidence-based strategy to support breastfeeding, particularly in settings with limited access to advanced lactation support. With growing evidence from both clinical trials and physiological studies, oxytocin massage holds promise as a standard non-pharmacological intervention in postpartum care, aligning with global efforts to improve maternal and infant health outcomes through sustainable breastfeeding practices.

This study has several limitations that should be considered when interpreting the findings. First, the study involved a relatively small sample size from a single public health center, which may limit the generalizability of the results to other populations and settings. Second, the follow-up period was limited to three days after the intervention, preventing the assessment of long-term effects of oxytocin massage on breast milk volume and infant growth. Third, although randomization was performed, potential confounding factors affecting breastfeeding outcomes, such as maternal nutritional status, psychological condition, family support, and breastfeeding frequency, were not fully controlled. In addition, age matching between participants was not conducted, and breast milk quality was not assessed.

Despite these limitations, the study has several strengths. The randomized single-blind controlled trial design reduced selection and assessment bias. Standardized intervention procedures were implemented by trained therapists, and outcome measurements were conducted using calibrated instruments and uniform protocols. Furthermore, the study evaluated both breast milk volume and infant weight outcomes, providing a more comprehensive assessment of the effectiveness of oxytocin massage in early postpartum women.

Future studies are warranted to investigate the long-term effects of oxytocin massage on exclusive breastfeeding rates, its impact on maternal stress and bonding, and comparisons with other lactation-enhancing interventions. Additionally, large-scale trials involving diverse populations could strengthen the generalizability of these findings and guide the development of standardized protocols for clinical practice.

Conclusion

A 30-minute oxytocin massage may be a potential supportive intervention for improving breast milk volume among women in the early postpartum period. The findings suggest that this intervention may also be associated with short-term improvements in infant weight gain. These findings highlight the potential value of implementing this protocol among mothers in the early postpartum period to support breastfeeding and promote early infant health.

Declarations

Acknowledgments

The authors would like to thank all research participants, including postpartum mothers, midwives, and the Head of Baregbeg Community Health Center, as well as the Institute for Research and Community Service (LPPM) of Universitas Galuh, for supporting and facilitating this study.

Conflict Of Interest

The authors declared no conflicts of interest.

Ethical Considerations

Written informed consent was obtained from all participants, and data were collected anonymously. Participants were informed about the study objectives, their right to decline responding to the questionnaire, and their freedom to discontinue or withdraw from the study at any time.

Code Of Ethics

All study procedures received ethical approval from the Dian Nuswantoro University HREC under protocol approval 319/EA/KEPK-Fkes-UDINUS/XI/2022

Use Of Artificial Intelligence (AI)

Not applicable.

Funding

This study was supported by Research Fund No. 037/SP2H/LT-MONO/LL4/2020 from the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia (RISTEKDIKTI).

Authors' Contributions

KDP and YIH contributed substantially to the study conception and design and were responsible for data collection. EYK and IP performed the data analysis and interpretation. The initial manuscript draft was prepared by KDP and EYK. IP also contributed to the study design, interpretation of findings, and critical revision of the manuscript for important intellectual content. All authors reviewed and approved the final version of the manuscript and accept responsibility for all aspects of the work, ensuring that any issues related to accuracy or integrity are appropriately addressed.

REFERENCES

1. World Health Organization. Breastfeeding [Internet]. 2023 [cited 2023 Dec 1]. Available from: https://www.who.int/health-topics/breastfeeding#tab=tab_1
2. The United Nations Children's Fund. World Breastfeeding Week 2021: Greater support needed for breastfeeding mothers in Indonesia amid COVID-19 [Internet]. 2021 [cited 2021 Dec 23]. Available from: <https://www.unicef.org/indonesia/press-releases/world-breastfeeding-week-2021-greater-support-needed-breastfeeding-mothers-indonesia>
3. World Health Organization. Global breastfeeding scorecard 2022: protecting breastfeeding through further investments and policy actions. World Health Organization; 2022.
4. World Health Organization. Exclusive breastfeeding for six months best for babies everywhere [Internet]. 2011 [cited 2021 Dec 23]. Available from: <https://www.who.int/news/item/15-01-2011-exclusive-breastfeeding-for-six-months-best-for-babies-everywhere>
5. World Health Organization. Infant and Young Child Feeding [Internet]. 2021 [cited 2022 Feb 4]. Available from: <https://www.who.int/news-room/fact-sheets/detail/infant-and-young-child-feeding>
6. Jabar D. Profil Kesehatan Jawa Barat Tahun 2019. Bandung: Dinas Kesehatan Jawa Barat; 2019. 61 p.
7. Ciamis D. Profil Kesehatan Kabupaten Ciamis Tahun 2019. Vol. 1. Ciamis: Dinas Kesehatan Kabupaten Ciamis; 2019. 71–74 p.
8. Nurdiana A, Siauta JA, Listiana R. Successful Pregnancy and Vaginal Delivery after Oxytocin Massage for Nulliparous Women with Anemia and Undetected Foetal

- Congenital Anomaly: An Indonesian Case Report. *EAS Journal of Nursing and Midwifery*. 2024;6(1). doi:10.36349/easjnm.2024.v06i01.002
9. Zahra S, Monireh A, Easa M, Susan P. Successful breastfeeding mothers' experiences of the difficulties of exclusive breastfeeding. *Acta Medica Mediterranea*[Internet]. 2015;31(7):1479–87.
10. Mishuk MH. Exploring Mother's Perception Regarding Exclusive Breastfeeding [Thesis] [Internet]. [Bangladesh]: Brac University; 2024 [cited 2026 Aug 16]. Available from: <https://dspace.bracu.ac.bd/items/8c9d63c2-a245-442f-af16-0e6744d9de6a>
11. Engelhart A, Mason S, Nwaozuru U, Obiezu-Umeh C, Carter V, Shato T, et al. Sustainability of breastfeeding interventions to reduce child mortality rates in low, middle-income countries: a systematic review of randomized controlled trials. *Frontiers in Health Services*. 2022;2:889390. doi:10.3389/frhs.2022.889390
12. Lestari P, Fatimah, Ayuningrum LD. The Effect of Oxytocin Massage During Postpartum on Baby Weight. *Jurnal Ners & Kebidanan Indonesia (Indonesian Journal of Nursing & Midwifery)*. 2021;9(2):147–54.
13. Lestari I, Rahmawati I, Emik Windarti H. SPEOS (Stimulation of Endorphin, Oxytocin and Suggestive): Intervention to Improvement of Breastfeeding Production. *Medico Legal Update*. 2019;19(1):210–5. doi:10.5958/0974-1283.2019.00042.2
14. Hesti KY, Pramono N, Wahyuni S, Widyawati MN, Santoso B. Effect of combination of breast care and oxytocin massage on breast milk secretion in postpartum mothers. *Belitung Nurs J*. 2017;3(6):784–90. doi:10.33546/bnj.293
15. UNICEF. Support Breastfeeding for A Healthier Planet [Internet]. 2020 [cited 2021 Dec 23]. Available from: <https://www.unicef.org/press-releases/support-breastfeeding-healthier-planet>
16. World Health Organization. Increasing commitment to breastfeeding through funding and improved policies and programmes: global breastfeeding scorecard 2019 [Internet]. 2019. Available from: <https://apps.who.int/iris/handle/10665/326049>
17. Hanieh S, Ha TT, Simpson JA, Thuy TT, Khuong NC, Thoang DD, et al. Exclusive breast feeding in early infancy reduces the risk of inpatient admission for diarrhea and suspected pneumonia in rural Vietnam: a prospective cohort study. *BMC Public Health*. 2015;15(1):1–10.
18. Triansyah A, Indarty A, Tahir M, Sabir M, Nur R, Basir-Cyio M, et al. The effect of oxytocin massage and breast care on the increased production of breast milk of breastfeeding mothers in the working area of the public health center of Lawanga of Poso District. *Gac Sanit*. 2021;35:S168–70. doi:10.1016/j.gaceta.2021.10.017
19. Verduci E, Gianni ML, Vizzari G, Vizzuso S, Cerasani J, Mosca F, et al. The triad mother-breast Milk-infant as predictor of future health: a narrative review. *Nutrients*. 2021;13(2):486. doi:10.3390/nu13020486
20. Boudry G, Charton E, Huerou-Luron L, Ferret-Bernard S, Le Gall S, Even S, et al. The relationship between breast milk components and the infant gut microbiota. *Front Nutr*. 2021;8:87. doi:10.3389/fnut.2021.629740
21. Kalyesubula M, Casey TM, Reicher N, Sabastian C, Wein Y, Shira EB, et al. Physiological state and photoperiod exposures differentially influence circadian rhythms of body temperature and prolactin and relate to changes in mammary PER1

- expression in late pregnant and early lactation dairy goats. *Small Ruminant Research*. 2021;200:106394. doi:10.1016/j.smallrumres.2021.106394
22. Dağlı E, Çelik N. The effect of oxytocin massage and music on breast milk production and anxiety level of the mothers of premature infants who are in the neonatal intensive care unit: A self-controlled trial. *Health Care Women Int*. 2022 May;43(5):1–14. doi:10.1080/07399332.2021.1947286
23. Johnson A, Kirk R, Rosenblum KL, Muzik M. Enhancing breastfeeding rates among African American women: A systematic review of current psychosocial interventions. *Breastfeeding Medicine*. 2014 Nov 25;10(1):45–62. doi:10.1089/bfm.2014.0023
24. Lebedevs T, Kendrick C. Pharmacological management of common lactation problems. *Journal of Pharmacy Practice and Research*. 2019;49(2):192–8. doi:10.1002/jppr.1561
25. Rodriguez IG, Leon FL, Corrales IG, Gonzalez JD. Impact and effectiveness of group strategies for supporting breastfeeding after birth: A systematic review. *Int J Environ Res Public Health*. 2021;18(5):2550. doi:10.3390/ijerph18052550
26. Kohan S, Keshvari M, Mohammadi F, Heidari Z. Designing and evaluating an empowering program for breastfeeding: a mixed-methods study. *Arch Iran Med*. 2019;22(8):443–52.
27. Umar. *Multitasking Breastfeeding Mama*. Jakarta: Pustaka Bunda Group; 2014.
28. Johan I, Azizah N. The Effect of Oxytocin Massage on Breastmilk Production Postpartum Mothers in Peterongan PHC Area, Jombang, East Java, Indonesia. *Global Nursing Challenges in The Free Trade Era* 5. 2016;1:1–9.
29. Dwedar LM, Osman HA, Mohamed HA, Ebrahim MM. Effect of Oxytocin versus Oketani Massage on Breastfeeding Predictors among Primipara Women. *Tanta Scientific Nursing Journal*. 2023;30(3):185–202. doi:10.21608/tsnj.2023.315147
30. Hendriyani F, Suryaningsih SA. Effectiveness of oxytocin massage and breast care treatment on breast milk production. *Indian J Forensic Med Toxicol*. 2019;13(4):4725–32.
31. Purnamasari KD, Hindiarti YI. Metode Pijat Oksitosin, Salah Satu Upaya Meningkatkan Produksi ASI Pada Ibu Postpartum. *Jurnal Kesehatan Perintis*. 2020;7(2):1–8.
32. Lakens D. Sample size justification. *Collabra Psychol*. 2022;8(1):33267.
33. Zlotnick C, Tzilos G, Miller I, Seifer R, Stout R. Randomized controlled trial to prevent postpartum depression in mothers on public assistance. *J Affect Disord*. 2016;189:263–8.
34. Damanik LPU. The effectiveness of neck massage in increasing puerperal mothers' breast milk quantity from day one to day three in Bantul. *Health Science Journal*. 2018;12(6):1–4. doi:10.21767/1791-809X.1000599
35. Goker A, Kosova F, Durgun SK, Demirtas Z. The effect of postpartum period back massage on serum ghrelin, leptin, adiponectin and visfatin levels. *Complement Ther Clin Pract*. 2021;42:101284. doi:10.1016/j.ctcp.2020.101284
36. Dini PR, Suwondo A, Hardjanti TR, Hadisaputro S, Mardiyono WMN. The Effect of Hypnobreastfeeding and Oxytocin Massage on Breastmilk Production in Postpartum. *Journal of Medical Science and Clinical Research*. 2017;5(10):28600–4.
37. Morhenn V, Beavin LE, Zak PJ. Massage increases oxytocin and reduces adrenocorticotropin hormone in humans. *Altern Ther Health Med*. 2012;18(6):11.

38. Swaab DF, Kreier F, Lucassen PJ, Salehi A, Buijs RM. Vasopressin and oxytocin beyond the pituitary in the human brain. *The Human Hypothalamus: Middle and Posterior Region*. 2021;180(3rd Series):7. doi:<https://doi.org/10.1016/B978-0-12-820107-7.00002-1>
39. Ly AD, Ningrum AG. Factor analysis of exclusive breastfeeding to mothers who work as health personnel in Kupang, Indonesia. *GSC Advanced Research and Reviews*. 2024;18(1):16–21. doi:10.30574/gscarr.2024.18.1.0491
40. Flaherman VJ, Murungi J, Bale C, Dickinson S, Chen X, Namiiro F, et al. Breastfeeding and Once-Daily Small-Volume Formula Supplementation to Prevent Infant Growth Impairment. *Pediatrics*. 2024;153(1):e2023062228. doi:10.1542/peds.2023-062228
41. Gustirini R, Anggraini IA. Combination of breast care and oxytocin massage of breastfeeding mothers in infant weight gain. *Jurnal Kesehatan Prima*. 2020;14(1):24–30. doi:10.32807/jkp.v14i1.287
42. Pertami SB, Budiono B, Rahmawati I. Optimizing the Endorphin and Oxytocin Massage to Increase Breast Milk Production among Postpartum Mother in Indonesia. *NurseLine Journal*. 2020;5(1):214–9. doi:10.19184/nlj.v5i1.16825
43. Lestari P, Fatimah F, Ayuningrum L, Herawati HD, Afifaturrohman N. Influence oxytocin massage on reduce lactation problems and support infants growth. *Open Access Maced J Med Sci*. 2022;10(T8):81–5. doi:10.3889/oamjms.2022.9487
44. El-Shamy FF, Ibrahim SM, Selim HK, EL Shwaikh SL, Salem SE, Elhosary EA. Ameliorative Potential of Relaxing Back Massage on Postpartum Women's Hormones and Infant Weight. *International Journal of Multiphysics*. 2025;19(1).